

Standard vaccination checklist (children 0-5 years of age)

Child's name:

Date of birth:

Age	Vaccine	Covered by National and/or School Immunisation Program	Tick
Birth	<ul style="list-style-type: none"> Hepatitis B 	Yes	<input type="checkbox"/>
6 weeks – 2 months	<ul style="list-style-type: none"> Hepatitis B, diphtheria, tetanus, whooping cough, haemophilus influenza, polio 	Yes	<input type="checkbox"/>
	<ul style="list-style-type: none"> Pneumococcal Rotavirus 	Yes Yes	<input type="checkbox"/> <input type="checkbox"/>
4 months	<ul style="list-style-type: none"> Hepatitis B, diphtheria, tetanus, whooping cough, haemophilus influenza, polio 	Yes	<input type="checkbox"/>
	<ul style="list-style-type: none"> Pneumococcal Rotavirus 	Yes Yes	<input type="checkbox"/> <input type="checkbox"/>
6 months	<ul style="list-style-type: none"> Hepatitis B, diphtheria, tetanus, whooping cough, haemophilus influenza, polio 	Yes	<input type="checkbox"/>
	<ul style="list-style-type: none"> Pneumococcal 	Yes	<input type="checkbox"/>
(6 months and up - annually)	<ul style="list-style-type: none"> Influenza 	<i>At-risk ATSI children; medically at-risk children</i>	<input type="checkbox"/>
12 months	<ul style="list-style-type: none"> Haemophilus influenza Meningococcal A, C, W, Y, tetanus Measles, mumps, rubella 	Yes Yes Yes	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
	<ul style="list-style-type: none"> Hepatitis A Pneumococcal (12-18 months) 	<i>At-risk ATSI persons</i> <i>At-risk ATSI children; medically at-risk children</i>	<input type="checkbox"/> <input type="checkbox"/>
	<ul style="list-style-type: none"> Meningococcal B* 	No	<input type="checkbox"/>
14 months	<ul style="list-style-type: none"> Meningococcal B (if given at 12 months) 	No	<input type="checkbox"/>
18 months	<ul style="list-style-type: none"> Diphtheria, tetanus, whooping cough, polio Measles, mumps, rubella, chickenpox 	Yes Yes	<input type="checkbox"/> <input type="checkbox"/>
	<ul style="list-style-type: none"> Hepatitis A 	<i>At risk ATSI persons</i>	<input type="checkbox"/>
4 years	<ul style="list-style-type: none"> Diphtheria, tetanus, whooping cough, polio 	Yes	<input type="checkbox"/>
	<ul style="list-style-type: none"> Pneumococcal 	<i>Medically at-risk children</i>	<input type="checkbox"/>

Please note: All children up to 19 years old can schedule 'catch-up' NIP vaccines for free if they did not receive it during childhood.

* The meningococcal B vaccine can be given from the age of 6 weeks and up. For those between the age of 6 weeks and 6 months, 3 doses are required (plus booster at 12 months) with intervals of 8 weeks in between. For those 6-12 months, 2 doses are required (plus booster dose 8 weeks after last primary dose) with intervals of 8 weeks in between. For those ≥ 12 months, 2 doses are required (no booster) with intervals of 8 weeks in between.